

AD / Tauopathy



Tau Seeding Mouse Model

P301L Tau AAV injections in the entorhinal cortex (ERC) of $\mathsf{APP}_{\mathsf{SI}}$ mice cause spreading of Tau in the hippocampus.


- · High total tau expression
- · Increased ptau Thr212/Ser214, Thr231, Ser262, Ser356, Ser396 levels
- · Increased AB levels
- · Pre-neurofibrillary tangles


Figure 1: Recombinant Adeno-associated virus serotype 9 (AAV9) with human P301L tau gene or empty vector was intracerebrally injected into the ERC of 3 months old male APP mice. Surgery coordinates: AP: -1.8; ML: +-1.4; DV: -1.4. A: Verification of injection site by ink injection into the entorhinal cortex. **B:** tau expression in the ERC after P301L tau virus injection. C: Spreading of tau into the hippocampus after P301L tau injection into the ERC of APP_{SL} mice. All analyses were performed 1


month after virus injection.


Figure 2: Rapid spread of virally induced pathological tau 4 months after injection. From left to right: Series of coronally cut brain


sections to visualize tau spreding throughout the hippocampus.














Scantox Discovery

Scantox Group, HQ Hestehavevej 36A, Ejby DK - 4623 Lille Skensved clientservice@scantox.com www.scantox.com

+45 5686 1500

© Scantox A/S Scantox is a registered trademark of Scantox A/S.

